

*An acne medicine
that's made to be loved*



Acanya ®
(Clindamycin Phosphate and
Benzoyl Peroxide) Gel, 1.2%/2.5%
Ready-to-Use 50g Pump

The power to treat
your acne without
overpowering
your skin

Acanya ®
(Clindamycin Phosphate and
Benzoyl Peroxide) Gel, 1.2%/2.5%
Ready-to-Use 50g Pump

The once-daily, topical treatment for acne vulgaris in patients 12 years of age or older

Proven to work



Acne is caused by several factors. Acanya® Gel combines 2 powerful medications that work in different ways to help you achieve results you'll love.

In clinical studies, Acanya Gel was proven to treat blackheads, whiteheads, and inflamed acne on the face.

Easy to use

- Wash your face gently and pat skin dry.
- Pump 1 pea-sized amount of Acanya Gel onto your fingertip and dot it onto your chin, both cheeks, nose, and forehead. Spread the gel over your face and gently rub it in.
- Do not get Acanya Gel in your mouth, eyes, or nose, on your lips, or in cuts or open wounds.
- Wash your hands with soap and water.

Avoid getting Acanya Gel in your hair or on colored fabric. Acanya Gel may bleach hair or colored fabric.



Designed to minimize skin irritation

- Acanya® Gel contains no alcohol or detergents, which may cause stinging or burning.
- May be used under makeup and sunscreen.
- Fragrance-free Acanya Gel goes on smoothly, disappears quickly, and leaves no residue.

Please note:

- Protect Acanya Gel-treated skin from sunlight by using an SPF 15 or higher sunscreen. Avoid tanning beds and sun lamps.
- Don't wash your face more than 2 to 3 times a day and avoid scrubbing your skin—it may make your acne worse.

Power to please

Acanya® Gel goes on smoothly and disappears quickly. Plus, the aqueous-based gel is designed specifically to help minimize skin irritation. Acanya Gel is a prescription medication that starts working fast.

Indication and Important Safety Information: Acanya Gel is indicated for the topical treatment of acne vulgaris in patients 12 years of age or older. Do not use Acanya Gel if you are allergic to clindamycin, have Crohn's disease, ulcerative colitis or have developed inflammation of the colon (colitis) with past antibiotic use. Clindamycin, the antibiotic in Acanya Gel, may cause diarrhea. If you experience severe diarrhea, bloody diarrhea, or severe abdominal cramping stop using Acanya Gel immediately and contact your doctor. Clindamycin taken orally or through IV may result in severe colitis, which may result in death. If you develop allergic symptoms such as severe itching, swelling of the face, eyes, lips, tongue or throat, or trouble breathing, you should stop using Acanya Gel and contact your doctor immediately. Limit your time in sunlight after applying Acanya Gel and avoid using tanning beds or sun lamps. Acanya Gel is for topical use only. Avoid applying Acanya Gel in mouth, eyes, or nose or on lips. Side effects may include redness, scaling, itching, burning, and stinging.

Get instant savings!

Pay no more than \$25 when you fill your prescription for Acanya[®] Gel.
Details and downloadable instant savings card at www.AcanyaGel.com.

Before using Acanya Gel, please read the accompanying patient information and instructions for applying. Talk to your doctor immediately about any side effect that bothers you or that does not go away. To learn more about Acanya Gel, visit www.AcanyaGel.com.



© 2011 CORIA Laboratories, a division of Valeant
Pharmaceuticals North America LLC ACAN-0211-0003

Acanya  [®]
(Clindamycin Phosphate and
Benzoyl Peroxide) Gel, 1.2%/2.5%
Ready-to-Use 50g Pump